

Question Paper
MS : JAM 2023

Section A: Q.1 – Q.10 Carry ONE mark each.

Q.1

Let $M = \begin{pmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{pmatrix}$. If a non-zero vector $X = (x, y, z)^T \in \mathbb{R}^3$ satisfies

$M^6 X = X$, then a subspace of \mathbb{R}^3 that contains the vector X is

(A) $\{(x, y, z)^T \in \mathbb{R}^3 : x = 0, y + z = 0\}$

(B) $\{(x, y, z)^T \in \mathbb{R}^3 : y = 0, x + z = 0\}$

(C) $\{(x, y, z)^T \in \mathbb{R}^3 : z = 0, x + y = 0\}$

(D) $\{(x, y, z)^T \in \mathbb{R}^3 : x = 0, y - z = 0\}$

Q.2 Let $M = M_1 M_2$, where M_1 and M_2 are two 3×3 distinct matrices. Consider the following two statements:

- (I) The rows of M are linear combinations of rows of M_2 .
- (II) The columns of M are linear combinations of columns of M_1 .

Then,

- (A) only (I) is TRUE
- (B) only (II) is TRUE
- (C) both (I) and (II) are TRUE
- (D) neither (I) nor (II) is TRUE

Q.3 Let $X \sim F_{6,2}$ and $Y \sim F_{2,6}$. If $P(X \leq 2) = \frac{216}{343}$ and $P\left(Y \leq \frac{1}{2}\right) = \alpha$, then 686α equals

- (A) 246
- (B) 254
- (C) 260
- (D) 264

Q.4 Let $Y \sim F_{4,2}$. Then, $P(Y \leq 2)$ equals

- (A) 0.60
- (B) 0.62
- (C) 0.64
- (D) 0.66

Q.5

Let X_1, X_2, \dots be a sequence of *i. i. d.* random variables each having $U(0,1)$ distribution.

Let Y be a random variable having the distribution function G . Suppose that

$$\lim_{n \rightarrow \infty} P\left(\frac{X_1 + X_2 + \dots + X_n}{n} \leq x\right) = G(x), \quad \text{for all } x \in \mathbb{R}.$$

Then, $\text{Var}(Y)$ equals

- (A) $\frac{1}{12}$
- (B) $\frac{1}{32}$
- (C) $\frac{1}{48}$
- (D) $\frac{1}{64}$

Q.6 Let X_1, X_2, X_3 be a random sample from an $N(\theta, 1)$ distribution, where $\theta \in \mathbb{R}$ is an unknown parameter. Then, which one of the following conditional expectations does NOT depend on θ ?

(A) $E(X_1 + X_2 - X_3 \mid X_1 + X_2)$

(B) $E(X_1 + X_2 - X_3 \mid X_2 + X_3)$

(C) $E(X_1 + X_2 - X_3 \mid X_1 - X_3)$

(D) $E(X_1 + X_2 - X_3 \mid X_1 + X_2 + X_3)$

Q.7 For the function $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x, y) = 2x^2 - xy - 3y^2 - 3x + 7y,$$

the point $(1, 1)$ is

(A) a point of local maximum

(B) a point of local minimum

(C) a saddle point

(D) NOT a critical point

Q.8 Let E_1, E_2 and E_3 be three events such that

$$P(E_1 \cap E_2) = \frac{1}{4}, \quad P(E_1 \cap E_3) = P(E_2 \cap E_3) = \frac{1}{5} \quad \text{and} \quad P(E_1 \cap E_2 \cap E_3) = \frac{1}{6}.$$


Then, among the events E_1, E_2 and E_3 , the probability that at least two events occur, equals

(A) $\frac{17}{60}$

(B) $\frac{23}{60}$

(C) $\frac{19}{60}$

(D) $\frac{29}{60}$



Q.9 Let X be a continuous random variable such that $P(X \geq 0) = 1$ and $Var(X) < \infty$. Then, $E(X^2)$ is

(A) $2 \int_0^{\infty} x^2 P(X > x) dx$

(B) $\int_0^{\infty} x^2 P(X > x) dx$

(C) $2 \int_0^{\infty} x P(X > x) dx$

(D) $\int_0^{\infty} x P(X > x) dx$

Q.10 Let X be a random variable having a probability density function

$$f(x; \theta) = \begin{cases} (3 - \theta) x^{2-\theta}, & \text{if } 0 < x < 1, \\ 0, & \text{otherwise,} \end{cases}$$

where $\theta \in \{0, 1\}$. For testing the null hypothesis $H_0: \theta = 0$ against $H_1: \theta = 1$, the power of the most powerful test, at the level of significance $\alpha = 0.125$, equals

(A) 0.15

(B) 0.25

(C) 0.35

(D) 0.45

Section A: Q.11 – Q.30 Carry TWO marks each.

Q.11 Let X_1 and X_2 be *i. i. d.* random variables having the common probability density function

$$f(x) = \begin{cases} e^{-x}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

Define $X_{(1)} = \min\{X_1, X_2\}$ and $X_{(2)} = \max\{X_1, X_2\}$. Then, which one of the following statements is FALSE?

(A) $\frac{2X_{(1)}}{X_{(2)} - X_{(1)}} \sim F_{2,2}$

(B) $2(X_{(2)} - X_{(1)}) \sim \chi_2^2$

(C) $E(X_{(1)}) = \frac{1}{2}$

(D) $P(3X_{(1)} < X_{(2)}) = \frac{1}{3}$

Q.12 Let X and Y be random variables such that $X \sim N(1, 2)$ and $P\left(Y = \frac{x}{2} + 1\right) = 1$. Let $\alpha = \text{Cov}(X, Y)$, $\beta = E(Y)$ and $\gamma = \text{Var}(Y)$. Then, the value of $\alpha + 2\beta + 4\gamma$ equals

- (A) 5
- (B) 6
- (C) 7
- (D) 8

Q.13 A point (a, b) is chosen at random from the rectangular region $[0, 2] \times [0, 4]$. Then, the probability that the area of the region

$$R = \{(x, y) \in \mathbb{R} \times \mathbb{R} : bx + ay \leq ab, \quad x, y \geq 0\}$$

will be less than 2, equals

- (A) $\frac{1 + \ln 2}{4}$
- (B) $\frac{1 + \ln 2}{2}$
- (C) $\frac{2 + \ln 2}{4}$
- (D) $\frac{1 + 2 \ln 2}{4}$

Q.14 Let X_1, X_2, \dots be a sequence of independent random variables such that

$$P(X_i = i) = \frac{1}{4} \quad \text{and} \quad P(X_i = 2i) = \frac{3}{4}, \quad i = 1, 2, \dots$$

For some real constants c_1 and c_2 , suppose that

$$\frac{c_1}{\sqrt{n}} \sum_{i=1}^n \frac{X_i}{i} + c_2 \sqrt{n} \xrightarrow{d} Z \sim N(0,1), \quad \text{as } n \rightarrow \infty.$$

Then, the value of $\sqrt{3} (3c_1 + c_2)$ equals

- (A) 2
- (B) 3
- (C) 4
- (D) 5

Q.15 Let X_1, X_2, \dots be a sequence of *i. i. d.* random variables such that

$$P(X_1 = 0) = P(X_1 = 1) = P(X_1 = 2) = \frac{1}{3}.$$

Let $S_n = \frac{1}{n} \sum_{i=1}^n X_i$ and $T_n = \frac{1}{n} \sum_{i=1}^n X_i^2$, $n = 1, 2, \dots$. Suppose that

$$\alpha_1 = \lim_{n \rightarrow \infty} P\left(\left|S_n - \frac{1}{2}\right| < \frac{3}{4}\right), \quad \alpha_2 = \lim_{n \rightarrow \infty} P\left(\left|S_n - \frac{1}{3}\right| < 1\right),$$


$$\alpha_3 = \lim_{n \rightarrow \infty} P\left(\left|T_n - \frac{1}{3}\right| < \frac{3}{2}\right) \quad \text{and} \quad \alpha_4 = \lim_{n \rightarrow \infty} P\left(\left|T_n - \frac{2}{3}\right| < \frac{1}{2}\right).$$

Then, the value of $\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4$ equals

- (A) 6
- (B) 5
- (C) 4
- (D) 3

Q.16 For $x \in \mathbb{R}$, the curve $y = x^2$ intersects the curve $y = x \sin x + \cos x$ at exactly n points. Then, n equals

- (A) 1
- (B) 2
- (C) 4
- (D) 8



Q.17 Let (X, Y) be a random vector having the joint probability density function

$$f(x, y) = \begin{cases} \alpha |x|, & \text{if } x^2 \leq y \leq 2x^2, \quad -1 \leq x \leq 1, \\ 0, & \text{otherwise,} \end{cases}$$

where α is a positive constant. Then, $P(X > Y)$ equals

(A) $\frac{5}{48}$

(B) $\frac{7}{48}$

(C) $\frac{5}{24}$

(D) $\frac{7}{24}$

Q.18 Let X_1, X_2, X_3, X_4 be a random sample of size 4 from an $N(\theta, 1)$ distribution, where $\theta \in \mathbb{R}$ is an unknown parameter. Let $\bar{X} = \frac{1}{4} \sum_{i=1}^4 X_i$, $g(\theta) = \theta^2 + 2\theta$ and $L(\theta)$ be the Cramer-Rao lower bound on variance of unbiased estimators of $g(\theta)$. Then, which one of the following statements is FALSE?

(A) $L(\theta) = (1 + \theta)^2$

(B) $\bar{X} + e^{\bar{X}}$ is a sufficient statistic for θ

(C) $(1 + \bar{X})^2$ is the uniformly minimum variance unbiased estimator of $g(\theta)$

(D) $Var((1 + \bar{X})^2) \geq \frac{(1+\theta)^2}{2}$

Q.19 Let X_1, X_2, \dots, X_n be a random sample from a population having the probability density function

$$f(x; \mu) = \begin{cases} \frac{1}{2} e^{-\left(\frac{x-2\mu}{2}\right)}, & \text{if } x > 2\mu, \\ 0, & \text{otherwise,} \end{cases}$$

where $-\infty < \mu < \infty$. For estimating μ , consider estimators

$$T_1 = \frac{\bar{X} - 2}{2} \quad \text{and} \quad T_2 = \frac{nX_{(1)} - 2}{2n},$$

where $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ and $X_{(1)} = \min\{X_1, X_2, \dots, X_n\}$. Then, which one of the following statements is TRUE?

(A) T_1 is consistent but T_2 is NOT consistent

(B) T_2 is consistent but T_1 is NOT consistent

(C) Both T_1 and T_2 are consistent

(D) Neither T_1 nor T_2 is consistent

Q.20 Let X_1, X_2, \dots, X_n be a random sample from a $U\left(\theta + \frac{\sigma}{\sqrt{3}}, \theta + \sqrt{3}\sigma\right)$ distribution, where $\theta \in \mathbb{R}$ and $\sigma > 0$ are unknown parameters. Let $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ and $S = \sqrt{\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2}$. Let $\hat{\theta}$ and $\hat{\sigma}$ be the method of moment estimators of θ and σ , respectively. Then, which one of the following statements is FALSE?

(A) $\hat{\sigma} + \sqrt{3}\hat{\theta} = \sqrt{3}\bar{X} - 3S$

(B) $2\sqrt{3}\hat{\sigma} + \hat{\theta} = \bar{X} - 4\sqrt{3}S$

(C) $\sqrt{3}\hat{\sigma} + \hat{\theta} = \bar{X} + \sqrt{3}S$

(D) $\hat{\sigma} - \sqrt{3}\hat{\theta} = 9S - \sqrt{3}\bar{X}$

Q.21

Let (X, Y, Z) be a random vector having the joint probability density function

$$f(x, y, z) = \begin{cases} \frac{1}{2xy}, & \text{if } 0 < z < y < x < 1, \\ \frac{1}{2x^2}, & \text{if } 0 < z < x < y < 2x < 2, \\ 0, & \text{otherwise.} \end{cases}$$

Then, which one of the following statements is FALSE?

(A) $P(Z < Y < X) = \frac{1}{2}$

(B) $P(X < Y < Z) = 0$

(C) $E(\min\{X, Y\}) = \frac{1}{4}$

(D) $\text{Var}(Y \mid X = \frac{1}{2}) = \frac{1}{12}$

Q.22 Let X be a random variable such that the moment generating function of X exists in a neighborhood of zero and

$$E(X^n) = (-1)^n \frac{2}{5} + \frac{2^{n+1}}{5} + \frac{1}{5}, \quad n = 1, 2, 3, \dots$$


Then, $P\left(\left|X - \frac{1}{2}\right| > 1\right)$ equals

(A) $\frac{1}{5}$

(B) $\frac{2}{5}$

(C) $\frac{3}{5}$

(D) $\frac{4}{5}$



Q.23 Let X be a random variable having a probability mass function $p(x)$ which is positive only for non-negative integers. If

$$p(x+1) = \left(\frac{\ln 3}{x+1}\right)p(x), \quad x = 0, 1, 2, \dots,$$

then $\text{Var}(X)$ equals

- (A) $\ln 3$
- (B) $\ln 6$
- (C) $\ln 9$
- (D) $\ln 18$

Q.24 Let $\{a_n\}_{n \geq 1}$ be a sequence such that $a_1 = 1$ and $4a_{n+1} = \sqrt{45 + 16a_n}$, $n = 1, 2, 3, \dots$. Then, which one of the following statements is TRUE?

- (A) $\{a_n\}_{n \geq 1}$ is monotonically increasing and converges to $\frac{17}{8}$
- (B) $\{a_n\}_{n \geq 1}$ is monotonically increasing and converges to $\frac{9}{4}$
- (C) $\{a_n\}_{n \geq 1}$ is bounded above by $\frac{17}{8}$
- (D) $\sum_{n=1}^{\infty} a_n$ is convergent

Q.25 Let the series S and T be defined by

$$\sum_{n=0}^{\infty} \frac{2 \cdot 5 \cdot 8 \cdots (3n+2)}{1 \cdot 5 \cdot 9 \cdots (4n+1)} \quad \text{and} \quad \sum_{n=1}^{\infty} \left(1 + \frac{1}{n}\right)^{-n^2},$$

respectively. Then, which one of the following statements is TRUE?

- (A) S is convergent and T is divergent
- (B) S is divergent and T is convergent
- (C) Both S and T are convergent
- (D) Both S and T are divergent

Q.26 The volume of the region

$$R = \{(x, y, z) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R} : x^2 + y^2 \leq 4, 0 \leq z \leq 4 - y\}$$

is

- (A) $16\pi - 16$
- (B) 16π
- (C) 8π
- (D) $16\pi + 4$

Q.27 For real constants α and β , suppose that the system of linear equations

$$x + 2y + 3z = 6; \quad x + y + \alpha z = 3; \quad 2y + z = \beta,$$

has infinitely many solutions. Then, the value of $4\alpha + 3\beta$ equals

- (A) 18
- (B) 23
- (C) 28
- (D) 32

Q.28

Let x_1, x_2, x_3 and x_4 be observed values of a random sample from an $N(\theta, \sigma^2)$ distribution, where $\theta \in \mathbb{R}$ and $\sigma > 0$ are unknown parameters. Suppose that $\bar{x} = \frac{1}{4} \sum_{i=1}^4 x_i = 3.6$ and $\frac{1}{3} \sum_{i=1}^4 (x_i - \bar{x})^2 = 20.25$. For testing the null hypothesis $H_0 : \theta = 0$ against $H_1 : \theta \neq 0$, the p -value of the likelihood ratio test equals

- (A) 0.712
- (B) 0.208
- (C) 0.104
- (D) 0.052

Q.29 Let X and Y be jointly distributed random variables such that, for every fixed $\lambda > 0$, the conditional distribution of X given $Y = \lambda$ is the Poisson distribution with mean λ . If the distribution of Y is $Gamma\left(2, \frac{1}{2}\right)$, then the value of $P(X = 0) + P(X = 1)$ equals

(A) $\frac{7}{27}$

(B) $\frac{20}{27}$

(C) $\frac{8}{27}$

(D) $\frac{16}{27}$

Q.30 Among all the points on the sphere $x^2 + y^2 + z^2 = 24$, the point (α, β, γ) is closest to the point $(1, 2, -1)$. Then, the value of $\alpha + \beta + \gamma$ equals

(A) 4

(B) -4

(C) 2

(D) -2

Section B: Q.31 – Q.40 Carry TWO marks each.

Q.31 Let M be a 3×3 real matrix. If $P = M + M^T$ and $Q = M - M^T$, then which of the following statements is/are always TRUE?

- (A) $\det(P^2 Q^3) = 0$
- (B) $\text{trace}(Q + Q^2) = 0$
- (C) $X^T Q^2 X = 0$, for all $X \in \mathbb{R}^3$
- (D) $X^T P X = 2X^T M X$, for all $X \in \mathbb{R}^3$

JAM 2023

JAM 2023

JAM 2023

Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অধ্যয়ন প্রবেশ পরীক্ষা

Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অধ্যয়ন প্রবেশ পরীক্ষা

Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অধ্যয়ন প্রবেশ পরীক্ষা

Organizing Institute
Indian Institute of Technology Guwahati

Organizing Institute
Indian Institute of Technology Guwahati

Organizing Institute
Indian Institute of Technology Guwahati

Q.32 Let X_1, X_2, X_3 be *i.i.d.* random variables, each having the $N(0, 1)$ distribution. Then, which of the following statements is/are TRUE?

(A)
$$\frac{\sqrt{2} (X_1 - X_2)}{\sqrt{(X_1 + X_2)^2 + 2X_3^2}} \sim t_1$$

(B)
$$\frac{(X_1 + X_2)^2}{(X_1 - X_2)^2 + 2X_3^2} \sim F_{1,2}$$

(C)
$$E\left(\frac{X_1}{X_2^2 + X_3^2}\right) = 0$$

(D)
$$P(X_1 < X_2 + X_3) = \frac{1}{3}$$

JAM 2023

JAM 2023

JAM 2023

Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অধ্যয়ন প্রবেশ পরীক্ষা

Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অধ্যয়ন প্রবেশ পরীক্ষা

Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অধ্যয়ন প্রবেশ পরীক্ষা

Organizing Institute
Indian Institute of Technology Guwahati

Organizing Institute
Indian Institute of Technology Guwahati

Organizing Institute
Indian Institute of Technology Guwahati

Q.33 Let x_1, x_2, \dots, x_{10} be the observed values of a random sample of size 10 from an $N(\theta, \sigma^2)$ distribution, where $\theta \in \mathbb{R}$ and $\sigma > 0$ are unknown parameters. If

$$\bar{x} = \frac{1}{10} \sum_{i=1}^{10} x_i = 0 \quad \text{and} \quad s = \sqrt{\frac{1}{9} \sum_{i=1}^{10} (x_i - \bar{x})^2} = 2,$$

then based on the values of \bar{x} and s and using Student's t -distribution with 9 degrees of freedom, 90% confidence interval(s) for θ is/are

(A) $(-0.8746, \infty)$

(B) $(-0.8746, 0.8746)$

(C) $(-1.1587, 1.1587)$

(D) $(-\infty, 0.8746)$

Q.34 Let (X_1, X_2) be a random vector having the probability mass function

$$f(x_1, x_2) = \begin{cases} \frac{c}{x_1! x_2! (12 - x_1 - x_2)!}, & \text{if } x_1, x_2 \in \{0, 1, \dots, 12\}, x_1 + x_2 \leq 12, \\ 0, & \text{otherwise,} \end{cases}$$

where c is a real constant. Then, which of the following statements is/are TRUE?

(A) $E(X_1 + X_2) = 8$

(B) $Var(X_1 + X_2) = \frac{8}{3}$

(C) $Cov(X_1, X_2) = -\frac{5}{3}$

(D) $Var(X_1 + 2X_2) = 8$

Q.35 Let P be a 3×3 matrix having the eigenvalues 1, 1 and 2. Let $(1, -1, 2)^T$ be the only linearly independent eigenvector corresponding to the eigenvalue 1. If the adjoint of the matrix $2P$ is denoted by Q , then which of the following statements is/are TRUE?

- (A) $\text{trace}(Q) = 20$
- (B) $\det(Q) = 64$
- (C) $(2, -2, 4)^T$ is an eigenvector of the matrix Q
- (D) $Q^3 = 20Q^2 - 124Q + 256I_3$

JAM 2023
Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অসামি প্ৰবেশ পরীক্ষা

JAM 2023
Organizing Institute
Indian Institute of Technology Guwahati

JAM 2023
Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অসামি প্ৰবেশ পরীক্ষা

JAM 2023

Organizing Institute
Indian Institute of Technology Guwahati

JAM 2023
Joint Admission test for Masters
সংযুক্ত স্নাতকোত্তর অসামি প্ৰবেশ পরীক্ষা

JAM 2023

Organizing Institute
Indian Institute of Technology Guwahati

Q.36 Let $f: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by

$$f(x, y) = \begin{cases} \frac{xy(x+y)}{x^2+y^2}, & \text{if } (x, y) \neq (0, 0), \\ 0, & \text{if } (x, y) = (0, 0). \end{cases}$$

Then, which of the following statements is/are TRUE?

- (A) f is continuous on $\mathbb{R} \times \mathbb{R}$
- (B) The partial derivative of f with respect to y exists at $(0, 0)$, and is 0
- (C) The partial derivative of f with respect to x is continuous on $\mathbb{R} \times \mathbb{R}$
- (D) f is NOT differentiable at $(0, 0)$

Q.37 Let X and Y be *i.i.d.* random variables each having the $N(0, 1)$ distribution. Let $U = \frac{X}{Y}$ and $Z = |U|$. Then, which of the following statements is/are TRUE?

- (A) U has a Cauchy distribution
- (B) $E(Z^p) < \infty$, for some $p \geq 1$.
- (C) $E(e^{tZ})$ does not exist for all $t \in (-\infty, 0)$
- (D) $Z^2 \sim F_{1,1}$

Q.38 Which of the following is/are TRUE?

(A) $\int_0^1 \int_0^1 e^{\max\{x^2, y^2\}} dx dy = e - 1$

(B) $\int_0^1 \int_0^1 e^{\min\{x^2, y^2\}} dx dy = \int_0^1 e^{t^2} dt - (e - 1)$

(C) $\int_0^1 \int_0^1 e^{\max\{x^2, y^2\}} dx dy = 2 \int_0^1 \int_y^1 e^{x^2} dx dy$

(D) $\int_0^1 \int_0^1 e^{\min\{x^2, y^2\}} dx dy = 2 \int_0^1 \int_1^y e^{y^2} dx dy$

Q.39 Let X be a random variable having the probability density function

$$f(x) = \begin{cases} \frac{5}{x^6}, & \text{if } x > 1, \\ 0, & \text{otherwise.} \end{cases}$$

Then, which of the following statements is/are TRUE for the distribution of X ?

(A) The coefficient of variation is $\frac{4}{\sqrt{15}}$

(B) The first quartile is $\left(\frac{3}{4}\right)^{\frac{1}{5}}$

(C) The median is $(2)^{\frac{1}{5}}$

(D) The upper bound obtained by Chebyshev's inequality for $P\left(X \geq \frac{5}{2}\right)$ is $\frac{1}{15}$

Q.40 Based on 10 data points $(x_1, y_1), (x_2, y_2), \dots, (x_{10}, y_{10})$ on a variable (X, Y) , the simple regression lines of Y on X and X on Y are obtained as $2y - x = 8$ and $y - x = -3$, respectively. Let $\bar{x} = \frac{1}{10} \sum_{i=1}^{10} x_i$ and $\bar{y} = \frac{1}{10} \sum_{i=1}^{10} y_i$. Then, which of the following statements is/are TRUE?

(A) $\sum_{i=1}^{10} x_i = 140$

(B) $\sum_{i=1}^{10} y_i = 110$

(C)
$$\frac{\sum_{i=1}^{10} (x_i - \bar{x}) y_i}{\sqrt{(\sum_{i=1}^{10} (x_i - \bar{x})^2)(\sum_{i=1}^{10} (y_i - \bar{y})^2)}} = -\frac{1}{\sqrt{2}}$$

(D)
$$\frac{\sum_{i=1}^{10} (x_i - \bar{x})^2}{\sum_{i=1}^{10} (y_i - \bar{y})^2} = 2$$

Section C: Q.41 – Q.50 Carry ONE mark each.

Q.41 Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by $f(x) = x^2 - x$, $x \in \mathbb{R}$. Let $g: \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable function such that $g(x) = 0$ has exactly three distinct roots in the open interval $(0, 1)$. Let $h(x) = f(x)g(x)$, $x \in \mathbb{R}$, and h'' be the second order derivative of the function h . If n is the number of roots of $h''(x) = 0$ in $(0, 1)$, then the minimum possible value of n equals _____

Q.42 Let X_1, X_2, \dots be a sequence of *i. i. d.* random variables, each having the probability density function

$$f(x) = \begin{cases} \frac{x^2 e^{-x}}{2}, & \text{if } x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

For some real constants β , γ and k , suppose that

$$\lim_{n \rightarrow \infty} P\left(\frac{1}{n} \sum_{i=1}^n X_i \leq x\right) = \begin{cases} 0, & \text{if } x < \beta, \\ kx, & \text{if } \beta \leq x \leq \gamma, \\ k\gamma, & \text{if } x > \gamma. \end{cases}$$

Then, the value of $2\beta + 3\gamma + 6k$ equals _____

Q.43 Let α and β be real constants such that

$$\lim_{x \rightarrow 0^+} \frac{\int_0^x \left(\frac{\alpha t^2}{1+t^4} \right) dt}{\beta x - \sin x} = 1.$$

Then, the value of $\alpha + \beta$ equals _____

Q.44 Let X_1, X_2, \dots, X_{10} be a random sample from an $N(0, \sigma^2)$ distribution, where $\sigma > 0$ is an unknown parameter. For some real constant c , let $Y = \frac{c}{10} \sum_{i=1}^{10} |X_i|$ be an unbiased estimator of σ . Then, the value of c equals _____ (round off to two decimal places)

Q.45 Let X be a random variable having the probability density function

$$f(x) = \begin{cases} \frac{x}{2}, & \text{if } 0 < x < 2, \\ 0, & \text{otherwise.} \end{cases}$$

Then, $Var\left(\ln\left(\frac{2}{X}\right)\right)$ equals _____

Q.46 Let X_1, X_2, X_3 be *i. i. d.* random variables, each having the $N(2, 4)$ distribution. If

$$P(2X_1 - 3X_2 + 6X_3 > 17) = 1 - \Phi(\beta),$$

then β equals _____

Q.47 Let the probability mass function of a random variable X be given by

$$P(X = n) = \frac{k}{(n-1)n}, \quad n = 2, 3, \dots,$$

where k is a positive constant. Then, $P(X \geq 17 \mid X \geq 5)$ equals _____

Q.48 Let

$$S_n = \sum_{k=1}^n \frac{1 + k 2^k}{4^{k-1}}, \quad n = 1, 2, \dots$$

Then, $\lim_{n \rightarrow \infty} S_n$ equals _____ (round off to two decimal places)

Q.49 A box contains a certain number of balls out of which 80% are white, 15% are blue and 5% are red. All the balls of the same color are indistinguishable. Among all the white balls, $\alpha\%$ are marked defective, among all the blue balls, 6% are marked defective and among all the red balls, 9% are marked defective. A ball is chosen at random from the box. If the conditional probability that the chosen ball is white, given that it is defective, is 0.4, then α equals _____

Q.50 Let X_1, X_2 be a random sample from a distribution having a probability density function

$$f(x; \theta) = \begin{cases} \frac{1}{\theta} e^{-\frac{x}{\theta}}, & \text{if } x > 0, \\ 0, & \text{otherwise,} \end{cases}$$

where $\theta \in (0, \infty)$ is an unknown parameter. For testing the null hypothesis $H_0 : \theta = 1$ against $H_1 : \theta \neq 1$, consider a test that rejects H_0 for small observed values of the statistic $W = \frac{X_1 + X_2}{2}$. If the observed values of X_1 and X_2 are 0.25 and 0.75, respectively, then the p -value equals _____ (round off to two decimal places)

Section C: Q.51 – Q.60 Carry TWO marks each.

Q.51 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by

$$f(x) = x^2 \sin(x - 1) + x e^{(x-1)}, \quad x \in \mathbb{R}.$$

Then,

$$\lim_{n \rightarrow \infty} n \left(f\left(1 + \frac{1}{n}\right) + f\left(1 + \frac{2}{n}\right) + \cdots + f\left(1 + \frac{10}{n}\right) - 10 \right)$$

equals _____

Q.52 Let the random vector (X_1, X_2) follow the bivariate normal distribution with

$$E(X_1) = E(X_2) = 1, \quad \text{Var}(X_2) = 4 \text{Var}(X_1) = 4 \quad \text{and} \quad \text{Cov}(X_1, X_2) = 1.$$

Then, $\text{Var}\left(X_1 + X_2 \mid X_1 = \frac{1}{2}\right)$ equals _____

Q.53 If, for some $\alpha \in (0, \infty)$,

$$\int_0^{\infty} 2^{-x^2} dx = \alpha \sqrt{\pi},$$

then α equals _____ (round off to two decimal places)

Q.54 Let $x_1 = 2.1$, $x_2 = 4.2$, $x_3 = 5.8$ and $x_4 = 3.9$ be the observed values of a random sample X_1, X_2, X_3 and X_4 from a population having a probability density function

$$f(x; \theta) = \begin{cases} \frac{x}{\theta^2} e^{-\frac{x}{\theta}}, & \text{if } x > 0, \\ 0, & \text{otherwise,} \end{cases}$$

where $\theta \in (0, \infty)$ is an unknown parameter. Then, the maximum likelihood estimate of $\text{Var}(X_1)$ equals _____

Q.55 Let $x_1 = 2$, $x_2 = 5$ and $x_3 = 4$ be the observed values of a random sample from a population having a probability mass function

$$f(x; \theta) = \begin{cases} \theta(1 - \theta)^x, & \text{if } x = 0, 1, 2, \dots, \\ 0, & \text{otherwise,} \end{cases}$$

where $\theta \in (0, 1)$ is an unknown parameter. If $\hat{\tau}$ is the uniformly minimum variance unbiased estimate of θ^2 , then $156 \hat{\tau}$ equals _____

Q.56 Let X_1, X_2, \dots, X_5 be *i. i. d.* random variables, each having the $Bin\left(1, \frac{1}{2}\right)$ distribution.
 Let $K = X_1 + X_2 + \dots + X_5$ and

$$U = \begin{cases} 0, & \text{if } K = 0, \\ X_1 + X_2 + \dots + X_K, & \text{if } K = 1, 2, \dots, 5. \end{cases}$$

Then, $E(U)$ equals _____

Q.57 Let $X_1 \sim \text{Gamma}(1, 4)$, $X_2 \sim \text{Gamma}(2, 2)$ and $X_3 \sim \text{Gamma}(3, 4)$ be three independent random variables. If $Y = X_1 + 2X_2 + X_3$, then $E\left(\left(\frac{Y}{4}\right)^4\right)$ equals _____

Q.58 Let X_1, X_2 be a random sample from a $U(0, \theta)$ distribution, where $\theta > 0$ is an unknown parameter. For testing the null hypothesis $H_0 : \theta \in (0, 1] \cup [2, \infty)$ against $H_1 : \theta \in (1, 2)$, consider the critical region

$$R = \left\{ (x_1, x_2) \in \mathbb{R} \times \mathbb{R} : \frac{5}{4} < \max\{x_1, x_2\} < \frac{7}{4} \right\}.$$

Then, the size of the critical region equals _____

Q.59 Let X_1, X_2, \dots, X_5 be a random sample from a $Bin(1, \theta)$ distribution, where $\theta \in (0, 1)$ is an unknown parameter. For testing the null hypothesis $H_0 : \theta \leq 0.5$ against $H_1 : \theta > 0.5$, consider the two tests T_1 and T_2 defined as:

T_1 : Reject H_0 if, and only if, $\sum_{i=1}^5 X_i = 5$.

T_2 : Reject H_0 if, and only if, $\sum_{i=1}^5 X_i \geq 3$.

Let β_i be the probability of making Type-II error, at $\theta = \frac{2}{3}$, when the test T_i , $i = 1, 2$, is used. Then, the value of $\beta_1 + \beta_2$ equals _____ (round off to two decimal places)

Q.60 Let $X_1 \sim N(2, 1)$, $X_2 \sim N(-1, 4)$ and $X_3 \sim N(0, 1)$ be mutually independent random variables. Then, the probability that exactly two of these three random variables are less than 1, equals _____ (round off to two decimal places)